



ARNI (Angiotensin Receptor blocker / Neprilysin Inhibitors [Sacubutril/Valsartan]) Heart Failure Medication Initiation and Titration



**This is a guide to monitoring; increase monitoring may be required given patient's status and co-morbidities (ie. renal insufficiency)

Angiotensin Converting Enzyme Inhibitors (ACEI) / Angiotensin Receptor Blockers (ARBS)*

Heart Failure Medication Initiation and Titration




*This algorithm is intended for single agent (ACEI or ARB)

**This is a guide to monitoring; increase monitoring may be required given patient's status and co-morbidities (ie. renal insufficiency)

Beta - Blockers

Heart Failure Medication Initiation and Titration




*verapamil and diltiazem are contraindicated in systolic heart failure (EF < 40%)

**This is a guide to monitoring; increase monitoring may be required given patient's status and co-morbidities


Diuretics

Heart Failure Medication Initiation and Titration



*Common signs and symptoms include new and worsening shortness of breath, weight gain, HF cough, orthopnea, paroxysmal nocturnal dyspnea, peripheral edema, ascites
 **Weight gain: 2-3 lbs in 1-2 days or 5 lbs in 1 week
 **This is a guide to monitoring; increased monitoring may be required given patient's status and co-morbidities
 Consider IV Lasix if non responsive to oral Lasix and Metolazone

Mineralocorticoid Receptor Antagonists (MRA) Heart Failure Medication Initiation and Titration



**This is a guide to monitoring; increase monitoring may be required given patient's status and co-morbidities (ie. renal insufficiency)